

University of Groningen

**Distribution of the urease gene cluster among and urease activities of enterohemorrhagic Escherichia coli O157 isolates from humans**

Friedrich, Alexander W; Köck, Robin; Bielaszewska, Martina; Zhang, Wenlan; Karch, Helge; Mathys, Werner

*Published in:*  
Journal of Clinical Microbiology

*DOI:*  
[10.1128/JCM.43.2.546-550.2005](https://doi.org/10.1128/JCM.43.2.546-550.2005)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2005

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Friedrich, A. W., Köck, R., Bielaszewska, M., Zhang, W., Karch, H., & Mathys, W. (2005). Distribution of the urease gene cluster among and urease activities of enterohemorrhagic Escherichia coli O157 isolates from humans. *Journal of Clinical Microbiology*, 43(2), 546-50. <https://doi.org/10.1128/JCM.43.2.546-550.2005>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## Distribution of the Urease Gene Cluster among and Urease Activities of Enterohemorrhagic *Escherichia coli* O157 Isolates from Humans

Alexander W. Friedrich,\* Robin Köck, Martina Bielaszewska, Wenlan Zhang,  
Helge Karch, and Werner Mathys

Institut für Hygiene, Universitätsklinikum Münster, Münster, Germany

Received 23 August 2004/Returned for modification 29 September 2004/Accepted 17 October 2004

**Enterohemorrhagic *Escherichia coli* (EHEC) O157 strains belong to two closely related major groups, which are differentiated by their sorbitol fermentation phenotypes. Here we studied the conservation of urease genes and their expression in sorbitol-fermenting (SF) and non-SF EHEC O157 isolates. PCR targeting *ure* genes (*ureA*, *-B*, *-C*, *-D*, *-E*, *-F*, and *-G*) demonstrated that each of these genes was present in 58 of 59 EHEC O157:H7 isolates. In contrast, none of 82 SF EHEC O157:NM (nonmotile) isolates contained any of the *ure* genes. Hence, the absence of the urease genes distinguishes SF EHEC O157:NM strains from EHEC O157:H7, but this absence demonstrates that the urease genes are not useful genetic targets for the detection of EHEC strains, because SF EHEC O157:NM strains are missed by such a strategy. When examined for urease activity on Christensen agar and in the API 20E system, only one O157:H7 strain displayed urease activity and produced elevated levels of ammonia, which was subsequently confirmed by ammonia electrode measurement. Because the *ure* genes were absent from each of nine strains of *E. coli* O55:H7, the proposed progenitor of EHEC O157, we hypothesize that EHEC O157:H7 diverged from the evolutionary pathway at an early stage and then acquired the O islands carrying the *ure* gene cluster.**

The enzyme urease catalyzes the hydrolysis of urea to give ammonia and carbon dioxide, thereby providing an important nitrogen source for many bacterial species. In addition, urease can contribute to the virulence of several gram-negative bacteria and enhance their acid resistance (24). Ureolytic *Escherichia coli* strains are rarely found among clinical isolates. Recently, the completed genome sequence analysis of enterohemorrhagic *E. coli* (EHEC) O157:H7 strain EDL933 (27) and the Sakai O157:H7 outbreak strain (13) demonstrated two and one loci, respectively, consisting of seven urease genes (*ureD*, *-A*, *-B*, *-C*, *-E*, *-F*, and *-G*). Those genes are clustered in the same fashion as found in other members of the family of *Enterobacteriaceae*. In EHEC strain EDL933, the urease gene cluster is located within O islands OI 43 and OI 48 (27). Subsequent studies indicated that among various diarrheagenic *E. coli* strains from clinical sources, urease genes were associated with major EHEC groups O26, O111, and O157 (26) but were absent from diarrheagenic *E. coli* strains of other pathogroups, including enterotoxigenic *E. coli*, enteroaggregative *E. coli*, enteroinvasive *E. coli*, and enteropathogenic *E. coli*. Among EHEC, particularly strains of serogroup O157, which cause diarrhea, hemorrhagic colitis, and the hemolytic-uremic syndrome (HUS), are most important (31). Based on sorbitol fermentation and geographic distribution, EHEC O157 belong to two major groups (21). While non-sorbitol-fermenting (non-SF) EHEC O157:H7 strains occur worldwide (31), SF EHEC O157:NM strains have emerged, so far, as important pathogens in Germany (10, 12, 21) and have caused two large outbreaks of HUS (2, 28). Multilocus enzyme electrophoresis, multilocus sequence typing, and determination of bacteriophage

integration sites indicate that SF EHEC O157:NM strains are closely related to EHEC O157:H7 (21, 29). Both groups possess *rfb*<sub>O157</sub> and *fliC* (encoding the H7 antigen), and most contain the *stx*<sub>2</sub> gene. Here we compared the two EHEC O157 groups for the presence of *ure* genes and urease production, using a large collection of EHEC O157:H7 and SF EHEC O157:NM strains isolated from patients in Germany.

### MATERIALS AND METHODS

**Bacterial strains.** Fifty-nine EHEC O157:H7 and 82 SF EHEC O157:NM strains were isolated between 1987 and 2003 from patients in Germany, Austria, and the Czech Republic. The EHEC O157:NM strains were from 72 patients with HUS and 10 patients with diarrhea. The EHEC O157:H7 strains were isolated from 31 patients with HUS and 28 patients with diarrhea. Between 1987 and 1994, such strains were detected by PCR screening of stools for *stx* genes, using primers MK1 and MK2 (19), and subsequent colony blot hybridization of PCR-positive stool cultures with P<sup>32</sup>-labeled *stx*<sub>1</sub> and *stx*<sub>2</sub> probes (18). Between 1995 and 2003, EHEC O157 were isolated by immunomagnetic separation and culture on SMAC agar (20) followed by colony hybridization with digoxigenin-labeled *stx* probes (10) and agglutination of *stx*-positive colonies with anti-O157 serum (Oxoid, Unipath GmbH, Wesel, Germany). The nine *eae*-positive *E. coli* O55:H7 strains included strains TB156A and TB182A (5) and seven additional strains from our collection. The laboratory *E. coli* strain ATCC 25922 was included in the study as a negative control.

**Urease activity.** Cultures were examined for urease activity with Christensen urea agar plates and API 20E (BioMérieux, Marcy-l'Etoile, France) strips after overnight incubation at 37°C. The measurement of ammonia and the subsequent calculations were performed as described previously (7). Briefly, bacteria were cultured overnight at 37°C in 40 ml of urea broth supplemented with 1 μM NiCl<sub>2</sub> and dissolved into a 0.4 McFarland suspension. After a 20-min incubation at room temperature, 0.4 ml of 10 M NaOH was added, and the liberated ammonia was quantified with a 95–10 ammonia electrode (Orion Research, Cambridge, Mass.); ammonia concentrations were calculated by comparison with a standard curve. The procedure was performed three times per isolate to calculate the arithmetic mean and variances in measurement. Before use, the ammonia electrode was calibrated against a series of standard solutions. Differences between means of the ammonia concentration were calculated by performing the *t* test and Mann-Wilcoxon test. *P* values of <0.05 were considered to be statistically significant.

\* Corresponding author. Mailing address: Institut für Hygiene, Universitätsklinikum Münster, Robert-Koch Str. 41, 48149 Münster, Germany. Phone: 49-251-8355366. Fax: 49-251-8355341. E-mail: alexf@uni-muenster.de.

TABLE 1. PCR primers and conditions used in this study

Primer	Sequence	Target	PCR conditions			PCR product (bp)	Reference
			Denaturing	Annealing	Extension		
ureA-f ureA-r	5'-GAC TCC AAG AGA AAA AGA CAA ACT A-3' 5'-CAG ATT ATC GGA TTA TGG ACG GTA-3'	<i>ureA</i>	94°C, 60 s	63°C, 60 s	72°C, 40 s	271	This study
ureB-f ureB-r	5'-GCT ACC TGC AGT ATT ATC GTT GAA-3' 5'-ACC CAT TAT ATC TCC GCG GAA ACC G-3'	<i>ureB</i>	94°C, 60 s	63°C, 60 s	72°C, 60 s	166	This study
ureC-f ureC-r	5'-TCT AAC GCC ACA ACC TGT AC-3' 5'-GAG GAA GGC AGA ATA TTG GG-3'	<i>ureC</i>	94°C, 60 s	60°C, 60 s	72°C, 60 s	398	26
ureD-f ureD-r	5'-CGT CAT CAT GTC GGT CTG CTC A-3' 5'-GCG TGG CTC CGG CGT AGT TTT-3'	<i>ureD</i>	94°C, 60 s	63°C, 60 s	72°C, 40 s	569	This study
ureE-f ureE-r	5'-GAG ACC CCG GCT CAG ACA ACT-3' 5'-CGTGATTATGGGCGTGCACCT-3'	<i>ureE</i>	94°C, 60 s	63°C, 60 s	72°C, 60 s	400	This study
ureF-f ureF-r	5'-ACT GGA GTG GGC AGT GGA AGC-3' 5'-ACGGAATAATCGGGAATACTGGG-3'	<i>ureF</i>	94°C, 60 s	63°C, 60 s	72°C, 60 s	564	This study
ureG-f ureG-r	5'-GGT CCG GTC GGC TCA GGT AAA-3' 5'-GAT GTT TTG CAG ACC TTC ACC AC-3'	<i>ureG</i>	94°C, 60 s	63°C, 60 s	72°C, 60 s	518	This study

**PCR.** PCRs for detecting the *ure* genes (*ureA*, *-B*, *-C*, *-D*, *-E*, *-F*, and *-G*) were performed in a cycler from Biometra GmbH, Göttingen, Germany. PCR reagents were purchased from Peqlab Biotechnologie (Erlangen, Germany). The 50- $\mu$ l PCR mixture consisted of 5  $\mu$ l of bacterial suspension (ca.  $10^4$  bacteria), 5  $\mu$ l of 10-fold-concentrated polymerase synthesis buffer Y containing 2.0 mM  $MgCl_2$ , 10  $\mu$ l of Enhancer solution, 1  $\mu$ l of 10 mM deoxynucleoside triphosphates, 30 pmol of each primer, and 1.25 U of *Taq* DNA polymerase. PCR primers, conditions, and target genes are shown in Table 1.

**Southern blot hybridization.** Genomic DNAs from three reference strains (*E. coli* O157:H7 strain EDL933, SF *E. coli* O157:NM strain 493/89, and *E. coli* O55:H7 strain TB182A) were digested with HindIII and EcoRI (New England Biolabs, Frankfurt, Germany) for hybridization with *ureA*, *-B*, *-C*, *-D*, *-F*, and *-G* probes; for hybridization with the *ureE* probe, digestion was performed with HindIII only. DNA fragments were separated on a 0.8% (wt/vol) agarose gel, transferred to a nylon membrane (Roti-Nylon plus; Carl Roth GmbH, Karlsruhe, Germany) by capillary blotting, and fixed by UV cross-linking (CL-1000 UV cross-linker; UVP, Inc., Upland, Calif.). The membrane was hybridized with the probes that were derived from *E. coli* O157:H7 strain EDL933 by PCR with primers listed in Table 1 and labeled with digoxigenin-11-dUTP by using a DIG DNA labeling and detection kit (Roche Molecular Biochemicals, Mannheim, Germany). Stringent hybridization was achieved with the DIG DNA labeling and detection kit according to the manufacturer's instructions.

**Sequence analysis.** PCR products were purified (PCR purification kit; Qiagen, Hilden, Germany) and sequenced with an automated ABI Prism 3100 Avant Genetic Analyzer and the ABI Prism BigDye Terminator Ready Reaction Cycle Sequencing Kit (Perkin-Elmer Applied Biosystems, Weiterstadt, Germany). Sequences were analyzed with the DNASIS program (Hitachi Software) and compared with sequences from the National Center for Biotechnology Information database.

## RESULTS

**Distribution of *ure* genes among EHEC O157 and *E. coli* O55:H7 isolates.** Each of the seven *ure* genes was present in 58 of 59 EHEC O157:H7 strains isolated from patients with diarrhea or HUS (Table 2). In contrast, these genes were absent from each of 82 SF EHEC O157:NM strains and from each of nine *eae*-positive *E. coli* O55:H7 strains which did not contain *stx* genes.

**Southern blot hybridization.** To confirm the absence of the *ure* genes in SF EHEC O157:NM and *E. coli* O55:H7 strains, genomic DNAs from reference strains 493/89 and TB182A,

respectively, were hybridized with digoxigenin-labeled *ureA*, *-B*, *-C*, *-D*, *-E*, *-F*, and *-G* probes. *E. coli* O157:H7 strain EDL933 and *E. coli* strain ATCC 25922 were used as positive and negative controls, respectively. Southern blot analysis demonstrated that each of the seven *ure* probes hybridized to a single fragment of genomic DNA (of approximately 1.2 and 9 kb) from EDL933 but not to DNA from any of the other three strains.

**Sequence analysis of the *ure* PCR amplicons.** As shown in Table 3, the nucleotide sequences of the PCR products derived from the *ure* genes of six representative *E. coli* O157:H7 strains were 99 to 100% identical to each other and to those of the corresponding *ure* genes from strain EDL933 (GenBank accession number NC002655) and the Sakai strain (GenBank accession number AP002554).

**Expression of urease activity and ammonia production.** When grown on Christensen agar and tested with API 20E strips, only one of the 150 strains investigated demonstrated urease activity. This strain (strain 4413/95) is of serotype O157:H7 and contains all of the *ure* genes. After 24 h of culture in urea broth, ammonia production by strain 4413/95 was significantly higher than that by strains EDL933, SF EHEC O157:NM 819/02, and ATCC 25922, which were all negative

TABLE 2. *ure* genes and ureolytic activity in EHEC O157 and *E. coli* O55:H7 strains

Serotype	No. of isolates	No. PCR positive for <i>ure</i> genes ( <i>ureA</i> , <i>-B</i> , <i>-C</i> , <i>-D</i> , <i>-E</i> , <i>-F</i> , and <i>-G</i> )	No. positive for ureolytic activity by:	
			Christensen agar	API 20E
O157:H7	59	58	1 <sup>a</sup>	1 <sup>a</sup>
SF O157:NM	82	0	0	0
<i>E. coli</i> O55:H7	9	0	0	0

<sup>a</sup> The strain that tested positive with the API 20E is identical to the strain that tested positive on Christensen agar.

TABLE 3. Results of PCRs with primers complementary to *ure* genes and nucleotide sequence homology of the PCR products to *ure* genes of EHEC O157:H7 strain EDL933

Strain no./yr of isolation	Serotype	Sorbitol fermentation	PCR results (% nucleotide sequence homology to EDL 933)						
			<i>ureA</i>	<i>ureB</i>	<i>ureC</i>	<i>ureD</i>	<i>ureE</i>	<i>ureF</i>	<i>ureG</i>
4413/95	O157:H7	—	+ (100)	+ (99)	+ (100)	+ (100)	+ (100)	+ (100)	+ (100)
HUS 1249	O157:H7	—	+ (100)	+ (100)	+ (99)	+ (100)	+ (100)	+ (99)	+ (100)
HC 3159	O157:H7	—	+ (100)	+ (100)	+ (100)	+ (100)	+ (100)	+ (100)	+ (99)
H631/88	O157:H7	—	+ (100)	+ (99)	+ (100)	+ (100)	+ (100)	+ (100)	+ (100)
H903-36/92	O157:H7	—	+ (100)	+ (99)	+ (100)	+ (100)	+ (100)	+ (100)	+ (100)
1194	O157:H7	—	+ (100)	+ (99)	+ (100)	+ (100)	+ (100)	+ (100)	+ (100)
5781/91	O157:H7	—	—	—	—	—	—	—	—
5869/96	O157:NM	+	—	—	—	—	—	—	—
3573/98	O157:NM	+	—	—	—	—	—	—	—
301/00	O157:NM	+	—	—	—	—	—	—	—
679/02	O157:NM	+	—	—	—	—	—	—	—
493/89	O157:NM	+	—	—	—	—	—	—	—
486/03	O55:H7	+	—	—	—	—	—	—	—
ATCC 25922		+	—	—	—	—	—	—	—

on Christensen agar (Fig. 1). Strains EDL933, 819/02, and ATCC 25922 produced comparably low levels of ammonia (Fig. 1).

# DISCUSSION

Since their first identification (17), SF EHEC strains of serotype O157:NM have emerged as important causes of diarrhea, bloody diarrhea, and HUS in Germany (10, 12, 21, 23). During the last several years they have been identified as human pathogens in other European and non-European countries, including Austria, the Czech Republic, Finland, Hungary, the United Kingdom, and Australia (1, 3, 4, 12, 21, 22). SF Shiga toxin-producing *E. coli* (STEC) O157:NM infections occur most frequently during cold months and disproportionately affect children under 3 years of age (12, 21). SF EHEC O157:NM strains are characterized by a specific combination of their phenotypic and virulence features, which differentiates them from non-SF EHEC O157:H7. Specifically, SF STEC O157:NM strains can ferment sorbitol overnight and produce  $\beta$ -D-glucuronidase; are assigned to phage type 88 or 23; possess *stx*<sub>2</sub> as their sole *stx* gene (21); do not contain the tellurite

resistance and adherence-conferring island (TAI) (30) but possess the complete EHEC factor-for-adherence (*efa1*) gene (15), which is only rudimentarily present in STEC O157:H7 (15); and also contain the *cdt* cluster, which is only rarely found in EHEC O157:H7 (16). Moreover, an *sfp* cluster encoding a putative adhesin has been identified on the large plasmid of SF EHEC O157:NM and has been shown to be characteristic of such strains (11). SF STEC O157:NM strains are nonmotile, but they regularly possess the *fliC* gene, encoding the H7 antigen (21). Monday et al. (25) have recently shown that the loss of motility in SF *E. coli* O157:NM strains is due to a 12-bp in-frame deletion in *fliH* that is required for transcriptional activation of genes involved in flagellum biosynthesis. By investigating the distribution of urease-encoding genes among EHEC O157 strains, we demonstrate in this study that EHEC O157:H7 and SF O157:NM differ remarkably from each other by the presence of the *ure* gene cluster.

The absence of *ure* genes from SF O157:NM strains, which has not been previously reported, has potential diagnostic implications. Nakano et al. (26) recently reported that the presence of *ureC* distinguishes EHEC of the major serogroups from diarrheagenic *E. coli* of other pathogroups and recommended this gene as a target to screen for EHEC. Our data clearly demonstrate that SF EHEC O157:NM strains would be missed by such an approach. On the other hand, PCR targeting *ureC* does provide a useful molecular tool for distinguishing SF EHEC O157:NM from EHEC O157:H7.

Multilocus enzyme electrophoresis demonstrates that SF EHEC O157:NM strains represent a distinct lineage within the EHEC O157:H7 clone complex (9, 34). In this evolutionary scenario, EHEC O157 evolved from an *E. coli* O55:H7 ancestor and then diverged into two groups, resulting in the O157:H7 clone, which spread worldwide, and the SF O157:NM clone, which is an increasing public health problem in Europe (9). Since the *ure* gene cluster is absent from both the *E. coli* O55:H7 strains and the SF EHEC O157:NM strains, it is possible that the EHEC O157:H7 strains have acquired the *ure* gene cluster later in their evolution by horizontal gene transfer.

Urease expression could modify internal and/or surrounding anion concentrations, enabling EHEC to survive acidic conditions and perhaps contributing to its low infectious dose and its




FIG. 1. Comparison of bacterial ammonia production after 24 h of culture in urea broth. Means  $\pm$  standard deviations of ammonia measurements for *E. coli* strain ATCC 25922, EHEC O157:H7 strain 4413/95, SF O157:NM strain 493/89, and EHEC O157:H7 strain EDL933 are shown. The data presented are from three independent experiments.



survival in the environment. In our study, only one of the 58 *ureC*-positive EHEC O157:H7 strains, which were isolated from patients with HUS or diarrhea, produced urease. The absence of urease production by the vast majority of EHEC O157 strains indicates that other acid tolerance mechanisms may be dominant in such strains. This low frequency of ureolytic activity in patients' isolates is similar to that found in *E. coli* O157:H7 strains isolated from patients in the United States (0.5%) (6) and Japan (7%) (35). In contrast, a substantially higher frequency of urease production (between 22 and 40%) has been reported in *E. coli* O157:H7 of bovine origin isolated in different countries (8, 32, 33). The reason for the low rate of urease expression among *E. coli* O157:H7 strains that are pathogenic for humans is unknown. Urease synthesis can be nitrogen regulated, urea inducible, or constitutive (24). Recent investigations demonstrated that the cloned EDL933 *ure* gene cluster was capable of synthesizing urease in an *E. coli* laboratory strain (14). However, when the *ure* gene cluster was transformed back into the native EDL933 background, the enzyme was not detectable, leading to speculation that an unidentified *trans*-acting factor(s) somehow down-regulates *ure* expression (14).

Because the *ure* gene cluster is regularly found in EHEC of serogroups O26, O111, and O157:H7, but not in other diarrheagenic *E. coli* strains (26), the question arises whether the presence of these genes contributes to the ability of EHEC to cause severe extraintestinal host injury. The fact that 88% of SF EHEC O157:NM strains in our study were isolated from HUS patients but none of these strains possessed urease-encoding genes suggests that the presence of these genes is not necessary for the ability of SF EHEC O157 to cause severe disease. However, if expressed *in vivo*, survival in the environment may be supported by the presence of *ure* genes in other EHEC strains.

#### ACKNOWLEDGMENTS

This study was supported by a grant from the Bundesministerium für Bildung und Forschung (BMBF) Project Network of Competence Pathogenomics Alliance "Functional Genomic Research on Enterohemorrhagic *Escherichia coli*" (BD number 119523).

We thank P. I. Tarr (Washington University School of Medicine, St. Louis, Mo.) for critical reading of the manuscript and fruitful discussions.

#### REFERENCES

- Alison, L. 2003. HUS due to a sorbitol-fermenting verotoxigenic *E. coli* O157 in Scotland. *Eurosurveillance Wkly.* 6:2-3.
- Ammon, A., L. R. Peterson, and H. Karch. 1999. A large outbreak of hemolytic uremic syndrome caused by an unusual sorbitol-fermenting strain of *E. coli* O157:H<sup>-</sup>. *J. Infect. Dis.* 179:1274-1277.
- Bettelheim, K. A., M. Whipp, S. P. Djordjevic, and V. Ramachandran. 2002. First isolation outside Europe of sorbitol-fermenting verocytotoxigenic *Escherichia coli* (VTEC) belonging to O group O157. *J. Med. Microbiol.* 51:713-714.
- Bielaszewska, M., H. Schmidt, M. A. Karmali, R. Khakhria, J. Janda, K. Blahova, and H. Karch. 1998. Isolation and characterization of sorbitol-fermenting Shiga toxin-producing *Escherichia coli* O157:H<sup>-</sup> strains in the Czech Republic. *J. Clin. Microbiol.* 36:2135-2137.
- Bokete, T. N., T. S. Whittam, R. A. Wilson, C. R. Clausen, C. M. O'Callahan, S. L. Moseley, T. R. Fritsche, and P. I. Tarr. 1997. Genetic and phenotypic analysis of *Escherichia coli* with enteropathogenic characteristics isolated from Seattle children. *J. Infect. Dis.* 175:1382-1389.
- Bopp, C. A., K. D. Greene, F. P. Downes, E. G. Sowers, J. G. Wells, and I. K. Wachsmuth. 1987. Unusual verotoxin-producing *Escherichia coli* associated with hemorrhagic colitis. *J. Clin. Microbiol.* 25:1486-1489.
- Butcher, G. P., S. D. Ryder, S. J. Hughes, M. Stewart, N. Bird, M. T. Haqqani, and J. M. Rhodes. 1992. Use of an ammonia electrode for rapid quantification of *Helicobacter pylori* urease: its use in the endoscopy room and in the assessment of urease inhibition by bismuth subsalicylate. *Digestion* 53:142-148.
- Conedera, G., S. Marangon, P. A. Chapman, A. Zuin, and A. Caprioli. 1997. Atypical strains of verocytotoxin-producing *Escherichia coli* O157 in beef cattle at slaughter in Veneto region, Italy. *Zentralbl. Veterinärmed. B* 44:301-306.
- Feng, P., K. A. Lampel, H. Karch, and T. S. Whittam. 1998. Genotypic and phenotypic changes in the emergence of *Escherichia coli* O157:H7. *J. Infect. Dis.* 177:1750-1753.
- Friedrich, A. W., M. Bielaszewska, W.-L. Zhang, M. Pulz, T. Kuczius, A. Ammon, and H. Karch. 2002. *Escherichia coli* harboring Shiga toxin 2 gene variants: frequency and association with clinical symptoms. *J. Infect. Dis.* 185:74-84.
- Friedrich, A. W., K. V. Nierhoff, M. Bielaszewska, A. Mellmann, and H. Karch. 2004. The pilin subunit gene (*sfpA*) of sorbitol-fermenting, enterohemorrhagic *Escherichia coli* O157:H<sup>-</sup>: phylogeny, clinical associations and diagnostic utility. *J. Clin. Microbiol.* 42:4697-4701.
- Gerber, A., H. Karch, F. Allerberger, H. M. Verwey, and L. B. Zimmerhackl. 2002. Clinical course and the role of Shiga toxin-producing *Escherichia coli* infection in the hemolytic-uremic syndrome in pediatric patients, 1997-2000, in Germany and Austria: a prospective study. *J. Infect. Dis.* 186:493-500.
- Hayashi, T., K. Makino, M. Ohnishi, K. Kurokawa, K. Ishii, K. Yokoyama, C. G. Han, E. Ohtsubo, K. Nakayama, T. Murata, M. Tanaka, T. Tobe, T. Iida, H. Takami, T. Honda, C. Sasakawa, N. Ogasawara, T. Yasunaga, S. Kuhara, T. Shiba, M. Hattori, and H. Shinagawa. 2001. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.* 8:11-22.
- Heimer, S. R., R. R. A. Welch, N. T. Perna, G. Posfai, P. S. Evans, J. B. Kaper, F. R. Blattner, and H. L. T. Mobley. 2002. Urease of enterohemorrhagic *Escherichia coli*: evidence for regulation by Fur and a *trans*-acting factor. *Infect. Immun.* 70:1027-1031.
- Janka, A., M. Bielaszewska, U. Dobrindt, and H. Karch. 2002. Identification and distribution of the enterohemorrhagic *Escherichia coli* factor for adherence (*efa1*) gene in sorbitol-fermenting *Escherichia coli* O157:H<sup>-</sup>. *Int. J. Med. Microbiol.* 292:207-214.
- Janka, A., M. Bielaszewska, U. Dobrindt, L. Greune, M. A. Schmidt, and H. Karch. 2003. The cytolethal distending toxin (*cdt*) gene cluster in enterohemorrhagic *Escherichia coli* O157:H<sup>-</sup> and O157:H7: characterization and evolutionary considerations. *Infect. Immun.* 71:3634-3638.
- Karch, H., R. Wiss, H. Glönnig, P. Emmrich, S. Aleksic, and J. Bockemühl. 1990. Hämolytisch-urämisches Syndrom bei Kleinkindern durch Verotoxin-produzierende *Escherichia coli*. *Dtsch. Med. Wochenschr.* 115:485-495.
- Karch, H., and T. Meyer. 1989. Evaluation of oligonucleotide probes for identification of Shiga-like-toxin-producing *Escherichia coli*. *J. Clin. Microbiol.* 27:1180-1186.
- Karch, H., and T. Meyer. 1989. Single primer pair for amplifying segments of distinct Shiga-like-toxin genes by polymerase chain reaction. *J. Clin. Microbiol.* 27:2751-2757.
- Karch, H., C. Janetzki-Mittmann, S. Aleksic, and M. Datz. 1996. Isolation of enterohemorrhagic *Escherichia coli* O157 strains from patients with hemolytic-uremic syndrome by using immunomagnetic separation, DNA-based methods, and direct culture. *J. Clin. Microbiol.* 34:516-519.
- Karch, H., and M. Bielaszewska. 2001. Sorbitol-fermenting Shiga toxin-producing *Escherichia coli* O157:H<sup>-</sup> strains: epidemiology, phenotypic and molecular characteristics, and microbiological diagnosis. *J. Clin. Microbiol.* 39:2043-2049.
- Keskimäki, M., M. Saari, T. Heiskanen, and A. Siitonen. 1998. Shiga toxin-producing *Escherichia coli* in Finland from 1990 through 1997: prevalence and characteristics of isolates. *J. Clin. Microbiol.* 36:3641-3646.
- Liesegang, A., U. Sachse, R. Prager, H. Claus, H. Steinrück, S. Aleksic, W. Rabsch, W. Voigt, A. Fruth, H. Karch, J. Bockemühl, and H. Tschäpe. 2000. Clonal diversity of Shiga toxin-producing *Escherichia coli* O157:H7/H<sup>-</sup> in Germany—a ten-year study. *Int. J. Med. Microbiol.* 290:269-278.
- Mobley, H. L., M. D. Island, and R. P. Hausinger. 1995. Molecular biology of microbial ureases. *Microbiol. Rev.* 59:451-480.
- Monday, S. R., S. A. Minnich, and P. C. Feng. 2004. A 12-base-pair deletion in the flagellar master control gene *flhC* causes nonmotility of the pathogenic German sorbitol-fermenting *Escherichia coli* O157:H<sup>-</sup> strains. *J. Bacteriol.* 186:2319-2327.
- Nakano, M., T. Iida, M. Ohnishi, K. Kurokawa, A. Takahashi, T. Tsukamoto, T. Yasunaga, T. Hayashi, and T. Honda. 2001. Association of the urease gene with enterohemorrhagic *Escherichia coli* strains irrespective of their serogroups. *J. Clin. Microbiol.* 39:4541-4543.
- Perna, N. T., G. Plunket III, V. Burland, B. Mau, J. D. Glasner, D. J. Rose, G. F. Mayhew, P. S. Evans, J. Gregor, H. A. Kirkpatrick, G. Posfai, J. Hackett, S. Klink, A. Boutin, Y. Shao, L. Miller, E. J. Grobeck, N. W. Davis, A. Lim, E. T. Dimalanta, K. D. Potamousis, J. Apodaca, T. S. Anantharaman, J. Lin, G. Yen, D. C. Schwartz, R. A. Welch, and F. R. Blattner. 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* 409:529-533.

28. **Robert Koch Institute.** 2002. Ein HUS-Ausbruch durch sorbitol-fermentierende EHEC des Serovars O157:H<sup>-</sup>: Untersuchungsergebnisse und Lehren für die Surveillance. *Epidemiologisches Bull.* **22**:171–175.
29. **Shaikh, N., and P. I. Tarr.** 2003. *Escherichia coli* O157:H7 Shiga toxin-encoding bacteriophages: integrations, excisions, truncations, and evolutionary implications. *J. Bacteriol.* **185**:3596–3605.
30. **Tarr, P. I., S. S. Bilge, J. C. Vary, S. Jelacic, R. L. Habeeb, T. R. Ward, M. R. Baylor, and T. E. Besser.** 2000. Iha: a novel *Escherichia coli* O157:H7 adherence-conferring molecule encoded on a recently acquired chromosomal island of conserved structure. *Infect. Immun.* **68**:1400–1407.
31. **Tarr, P. I., and M. A. Neill.** 2001. *Escherichia coli* O157:H7. *Gastroenterol. Clin. North Am.* **30**:735–751.
32. **Tutenel, A. V., D. Pierard, J. Uradzinski, E. Jozwik, M. Pastuszcak, J. Van Hende, M. Uyttendaele, J. Debevere, T. Cheasty, J. Van Hoof, and L. De Zutter.** 2002. Isolation and characterization of enterohaemorrhagic *Escherichia coli* O157:H7 from cattle in Belgium and Poland. *Epidemiol. Infect.* **129**:41–47.
33. **Tutenel, A. V., D. Pierard, J. Van Hoof, M. Cornelis, and L. De Zutter.** 2003. Isolation and molecular characterization of *Escherichia coli* O157 isolated from cattle, pigs and chickens at slaughter. *Int. J. Food Microbiol.* **84**:63–69.
34. **Whittam, T. S., M. L. Wolfe, I. K. Wachsmuth, F. Ørskov, I. Ørskov, and R. A. Wilson.** 1993. Clonal relationship among *Escherichia coli* strains that cause hemorrhagic colitis and infantile diarrhea. *Infect. Immun.* **61**:1619–1629.
35. **Yamada, F., T. Kurazono, M. Yamaguchi, Y. Ohzeki, and Y. Okuyama.** 1994. Biological characters of enterohemorrhagic *Escherichia coli* isolates from diarrhea patients in Saitama (1990–1992). *Kansenshogaku Zasshi* **68**:1451–1458.